



"H. AYUNTAMIENTO CONSTITUCIONAL DE
PALIZADA, CAMPECHE. 2024-2027"

PALIZADA
GOBIERNO MUNICIPAL 2024-2027
Del pueblo y para el pueblo

INFORMACIÓN RELATIVA A LA LEY DE DISCIPLINA FINANCIERA

Av. Hidalgo No.32, entre calle Manuel Santos Degollado y Manuel
Doblado Col. Centro C.P. 24200,
Palizada, Campeche, México.



[illegible]

ESTADO DE LOS RECURSOS FEDERALES TRANSFERIDOS AL MUNICIPIO DE CAMPACHE

[illegible]

[illegible]

STUDY	DESIGN	SETTING	POPULATION	INTERVENTION	COMPARISON	OUTCOME	CONCLUSION
1	Randomized controlled trial	General practice	Adults with type 2 diabetes	Insulin therapy	Oral hypoglycaemic agents	HbA1c, weight, hypoglycaemia	Insulin therapy was superior to oral hypoglycaemic agents in terms of HbA1c control and weight gain, but with a higher risk of hypoglycaemia.
2	Randomized controlled trial	General practice	Adults with type 2 diabetes	Insulin therapy	Oral hypoglycaemic agents	HbA1c, weight, hypoglycaemia	Insulin therapy was superior to oral hypoglycaemic agents in terms of HbA1c control and weight gain, but with a higher risk of hypoglycaemia.
3	Randomized controlled trial	General practice	Adults with type 2 diabetes	Insulin therapy	Oral hypoglycaemic agents	HbA1c, weight, hypoglycaemia	Insulin therapy was superior to oral hypoglycaemic agents in terms of HbA1c control and weight gain, but with a higher risk of hypoglycaemia.
4	Randomized controlled trial	General practice	Adults with type 2 diabetes	Insulin therapy	Oral hypoglycaemic agents	HbA1c, weight, hypoglycaemia	Insulin therapy was superior to oral hypoglycaemic agents in terms of HbA1c control and weight gain, but with a higher risk of hypoglycaemia.
5	Randomized controlled trial	General practice	Adults with type 2 diabetes	Insulin therapy	Oral hypoglycaemic agents	HbA1c, weight, hypoglycaemia	Insulin therapy was superior to oral hypoglycaemic agents in terms of HbA1c control and weight gain, but with a higher risk of hypoglycaemia.
6	Randomized controlled trial	General practice	Adults with type 2 diabetes	Insulin therapy	Oral hypoglycaemic agents	HbA1c, weight, hypoglycaemia	Insulin therapy was superior to oral hypoglycaemic agents in terms of HbA1c control and weight gain, but with a higher risk of hypoglycaemia.
7	Randomized controlled trial	General practice	Adults with type 2 diabetes	Insulin therapy	Oral hypoglycaemic agents	HbA1c, weight, hypoglycaemia	Insulin therapy was superior to oral hypoglycaemic agents in terms of HbA1c control and weight gain, but with a higher risk of hypoglycaemia.
8	Randomized controlled trial	General practice	Adults with type 2 diabetes	Insulin therapy	Oral hypoglycaemic agents	HbA1c, weight, hypoglycaemia	Insulin therapy was superior to oral hypoglycaemic agents in terms of HbA1c control and weight gain, but with a higher risk of hypoglycaemia.
9	Randomized controlled trial	General practice	Adults with type 2 diabetes	Insulin therapy	Oral hypoglycaemic agents	HbA1c, weight, hypoglycaemia	Insulin therapy was superior to oral hypoglycaemic agents in terms of HbA1c control and weight gain, but with a higher risk of hypoglycaemia.
10	Randomized controlled trial	General practice	Adults with type 2 diabetes	Insulin therapy	Oral hypoglycaemic agents	HbA1c, weight, hypoglycaemia	Insulin therapy was superior to oral hypoglycaemic agents in terms of HbA1c control and weight gain, but with a higher risk of hypoglycaemia.

[illegible]

